

Photon-photon interaction and Schwinger process

K. Bulycheva

*Institute for Theoretical and Experimental Physics, Moscow, Russia
Moscow Institute of Physics and Technology, Dolgoprudny, Russia
bulycheva@itep.ru*

S. Guts

*Institute for Theoretical and Experimental Physics, Moscow, Russia
Moscow Institute of Physics and Technology, Dolgoprudny, Russia
guts@itep.ru*

Abstract

We consider the tunneling processes induced by two-particle collisions. The result derived by the thermal bath approach is reproduced by the classical configurations. The Coulomb and Yukawa corrections for the rate of the process are computed.

1 Introduction

The problem of creation electron-positron pairs has been investigated since historical works of Schwinger and Euler-Heisenberg [1], [2]. However, this process hasn't been observed experimentally due to huge strengths of fields needed. The pair production rate is exponentially suppressed when the strength of E is lower than $\frac{m^2}{e}$, which is much larger than any available one. One can hope to obtain experimental evidence of the process by investigating energetic photons interacting with strong and low-frequency electrical field created by lasers [3], [4]. High-energy photons considerably enhance the rate of pair production. This phenomenon is called induced Schwinger process.

There are various types of Schwinger-like processes. They all are closely related to the tunnelling decay of false vacuum. We can investigate scalar field to obtain main phenomena arising in the problem [5], [7], [8], [10], [11]. These results can be generalized to the case of electron-positron pair creation [19]. Monopole decay in constant electrical field appears to be quite similar to induced Schwinger process [12], [15], [17], [18]. Schwinger type processes also can be investigated using the brane language [12], [13].

One of the natural generalizations of these processes is pair creation induced by two photons. This problem was investigated in different frameworks too [11], [20]. One of the most used technique to derive amplitudes of the process is introducing finite temperature [20].

In this paper the many-photon Schwinger process is investigated. The first subsection contains a direct calculation of the cross-section for two scalar particles interacting at the background of the bounce. This value was obtained in [20], and our technique is a cross-check of thermal bath approach. Also in this section the imaginary part of the correlator of arbitrarily many scalar fields at the background of the bounce is obtained. In the next section the polarization operator for two photons interacting at the background of the bounce is computed and the way of computing the polarization operator for many-photon case is shown. Finally, in the third subsection the results of the previous two subsections are used to find the Coulomb and Yukawa corrections to the width of the process. The Coulomb correction coincides with that calculated in [9].

2 Two-particle cross-sections

2.1 Scalar particle

To understand the structure of the cross-section, we consider the case of scalar field in two Euclidean dimensions and cross-check the result of [20]. Let's introduce two scalar fields ϕ and χ .

$$L = \frac{1}{2} (\partial_\mu \phi)^2 + \frac{1}{2} (\partial_\mu \chi)^2 - \frac{1}{2} m^2 \chi^2 - V(\phi) - V_{int}(\phi, \chi), \quad (1)$$

The potential $V(\phi)$ is quartic with "false" and "true" vacua:

$$V(\phi) = \frac{1}{4}\lambda(\phi^2 - \nu^2)^2 - \frac{\varepsilon}{2\nu}(\phi + \nu). \quad (2)$$

We use the limit when the mass of field χ is much smaller than the mass of ϕ .

$$m \ll M = \nu\sqrt{\lambda}, \quad (3)$$

We also need the condition of thin-wall approximation

$$R \gg 1/M, \quad (4)$$

where R is the radius of the bounce. This condition implies that the configuration being discussed is in quasi-classical regime, and quantum effects can be neglected. In this approximation the action can be divided to an internal part and a wall part. Minimizing the action, one can obtain the radius R .

$$S = S_{wall} + S_{int} = \mu L - \varepsilon A = 2\pi\mu R - \pi\epsilon R^2, \quad (5)$$

Here L is the perimeter and A is the area of the bubble. In the thin-wall limit $\mu = \nu^3 \frac{\sqrt{8\lambda}}{3}$. Then the radius of the bubble becomes $R = \frac{\mu}{\epsilon}$. The condition (4) becomes $\varepsilon \ll \lambda\nu^4$. The action then is

$$S = \frac{\pi\mu^2}{\epsilon}. \quad (6)$$

Interaction between the fields is given by

$$V_{int} = g\rho(\phi)\chi, \quad (7)$$

The exact form of interaction term is not very important, it must just be significantly different from zero when ϕ is far from vacuum. In the discussed approximation the support of ρ is very thin and coincides with the boundary of the circle and can be considered as δ -function.

$$\rho^B(x, t) = \frac{1}{2\pi R} \delta\left(\sqrt{x^2 + t^2} - R\right), \quad (8)$$

To compute the cross-section for process like $\vec{q}_1 + \vec{q}_2 \rightarrow \vec{q}_3 + \vec{q}_4$, one needs to follow [11] and calculate the imaginary part of the four-fold density correlator:

$$\Pi^{(4)}(x_1, t_1, x_2, t_2, x_3, t_3) = \langle \rho(t_1, x_1) \rho(t_2, x_2) \rho(t_3, x_3) \rho(0, 0) \rangle,$$




Figure 1: The bounce with four areas where the density of ϕ is concentrated. The centers of outside circles belong to board of the bold circle which refers to bounce. That implies that all the four outside circles intersect in a single point.

$$\text{Im } \Pi = \frac{\Gamma}{2L} \int \rho^B(x_1 - x, t_1 - t) \rho^B(x_2 - x, t_2 - t) \rho^B(x_3 - x, t_3 - t) \rho^B(x, t) dx dt. \quad (9)$$

Here $\frac{\Gamma}{L}$ is the rate of spontaneous vacuum decay. It was calculated in [6] and has the following form:

$$\frac{\Gamma}{2L} = \frac{\varepsilon}{4\pi} \exp \left\{ -\frac{\pi\mu^2}{\varepsilon} \right\}, \quad (10)$$

Geometrically, every ρ^B is a circle, and all four this circles need to lay on the board of initial bounce. This gives us additional condition on an relative location of (x_i, t_i) ; however, we shall see that this condition is implied automatically. Due to the congruence of radii of this circles with the radius of the initial bounce, this condition is equivalent to that of intersection of the four circles in one point, and this condition is provided by δ -functions (see fig. 1). Substituting (8) into (9), we obtain

$$\text{Im } \Pi = \frac{\Gamma}{2L} \frac{1}{(2\pi R)^4} \frac{4R^2}{\sqrt{x_1^2 + t_1^2} \sqrt{4R^2 - x_1^2 - t_1^2}} \delta \left(\sqrt{x_2^2 + t_2^2} - R \right) \delta \left(\sqrt{x_3^2 + t_3^2} - R \right). \quad (11)$$

To switch to the momentum representation instead of the coordinate one, we perform

Fourier transformation.

$$\begin{aligned}
& \int \frac{4R^2}{\sqrt{x_1^2 + t_1^2} \sqrt{4R^2 - x_1^2 - t_1^2}} \delta \left(\sqrt{x_2^2 + t_2^2} - R \right) \delta \left(\sqrt{x_3^2 + t_3^2} - R \right) \\
& \exp \{i(q_1, r_1) + i(q_2, r_2) + i(q_3, r_3)\} d^2 r_1 d^2 r_2 d^2 r_3 = \\
& -4\pi^2 R^2 I_0^2(mR) \int \delta(r_2 - R) \delta(r_3 - R) e^{i(q_2, r_2) + i(q_3, r_3)} d^2 r_2 d^2 r_3 = -16\pi^4 R^4 I_0^4(mR),
\end{aligned} \tag{12}$$

Here we use the fact that external momentum satisfies $|q_i| = m$. During this calculation the integral definitions of Bessel functions were used:

$$\begin{aligned}
& \int \frac{e^{i(q, r)}}{r \sqrt{4R^2 - r^2}} d^2 r = \int \frac{e^{iqr \cos \theta}}{\sqrt{4R^2 - r^2}} dr d\theta = - \int \frac{e^{2iqR \cos \phi \cos \theta}}{2R \sin \phi} 2R \sin \phi d\phi d\theta = \\
& = - \int e^{iqR(\cos(\phi - \theta) + \cos(\phi + \theta))} d\phi d\theta = -\frac{1}{2} \int e^{iqR(\cos \alpha + \cos \beta)} d\alpha d\beta = -\pi^2 I_0^2(mR),
\end{aligned} \tag{13}$$

The factor of $\frac{1}{2}$ arises due to the Jacobian of the transformation and disappears because of shift of the integration limits.

$$\int \delta(r - R) e^{i(q, r)} d^2 r = \int e^{iqR \cos \theta} R d\theta = 2\pi R J_0(qR) = 2\pi R I_0(mR), \tag{14}$$

Thus we come to exact expression for the cross-section:

$$\sigma = -\frac{\text{Im } \Pi}{2I} = \frac{\epsilon}{4\pi} \frac{1}{2I} (g I_0(mR))^4 \exp \left\{ -\frac{\pi \mu^2}{\epsilon} \right\}. \tag{15}$$

where $I = \sqrt{(q_1 q_2)^2 - m^4}$ is a kinematical invariant. One can make sure that this coincides with result (32) in [20] derived with thermal bath approach.

This result can be obtained by a quite simpler technique. As we have understood that geometric properties do not affect the answer, we can pass to the momentum representation and actually permute the order of integration in (11), (12). The density can be written in the momentum representation.

$$\int \rho^B(x_1 - x, t_1 - t) e^{i(q_1, r_1)} d^2 r = \int \rho^B(x_1, t_1) e^{i(q_1, r_1)} e^{i(q_1, r)} d^2 r = I_0(mR) e^{i(q_1, r)}, \tag{16}$$

Then the calculation transforms to computing four integrals of the same type:

$$\begin{aligned}
& \int \rho^B(x_1 - x, t_1 - t) \rho^B(x_2 - x, t_2 - t) \rho^B(x_3 - x, t_3 - t) \rho^B(x, t) \\
& \exp \{i(q_1, r_1) + i(q_2, r_2) + i(q_3, r_3)\} d^2 r_1 d^2 r_2 d^2 r_3 dx dt = \\
& \frac{1}{2\pi R} I_0^3(mR) \int \delta \left(\sqrt{x^2 + t^2} - R \right) e^{i(q_1 + q_2 + q_3, r)} d^2 r = -I_0^4(mR),
\end{aligned} \tag{17}$$

Due to momentum conservation law, $\vec{q}_1 + \vec{q}_2 + \vec{q}_3 = -\vec{q}_4$, $|q_4| = m$ in the last integral. From here one can see that an n -fold correlator can be expressed as

$$\text{Im } \Pi^{(n)} = -\frac{\Gamma}{2L} (gI_0(mR))^n. \quad (18)$$

2.2 Electron

The electron-positron pair in external electrical field in (1+1) Euclidean dimensions appears to have much in common with the case of the false vacuum decay. We can think of this process in terms of action (5), where μ refers to the mass of electron, ε arises due to screening the field by the charged pair, and the circular shape of the bounce corresponds to circular motion of the charge in Euclidean electrical field. Thus the technique of calculating the imaginary part of a four-fold correlator can help to find the polarization operator in leading order on $\frac{eE}{m^2}$ (neglecting the spinor structure of electron current). We take the electron current on the bounce in the form

$$j_B^\mu(x) = \frac{e}{2\pi R} \frac{n^\mu}{\sqrt{x^2 + t^2}} \delta(\sqrt{x^2 + t^2} - R) \delta(y) \delta(z), \quad (19)$$

where $n^\mu = (x, -t, 0, 0)^T = r(\sin \phi, -\cos \phi, 0, 0)^T$ stands for the unit tangential vector to the wall of the bounce. In this case the operator we need is

$$\text{Im } \Pi^{\mu\nu\kappa\rho} = \frac{\Gamma}{2V} \int j^\mu(x_1 - x) j^\nu(x_2 - x) j^\kappa(x_3 - x) j^\rho(-x) d^4x, \quad (20)$$

where prefactor $\frac{\Gamma}{2V}$ is computed, for example, in [9]:

$$\frac{\Gamma}{2V} = \frac{(eE)^2}{8\pi^3} \exp \left\{ -\frac{\pi m^2}{eE} \right\}. \quad (21)$$

To simplify the calculations and the shape of the answer, we switch to the momentum form (on (x, t) plane)).

$$j^\mu(q_1) = e^{i(q_1, x)} \int \frac{e}{2\pi R r} r \begin{pmatrix} \sin \phi \\ -\cos \phi \end{pmatrix} \delta(r - R) e^{iq_1 r \cos(\theta_1 - \phi)} r d\phi dr, \quad (22)$$

Here and in what follows, the q denotes the absolute value of the momentum in the (x, t) plane, i.e. $q = \sqrt{q_x^2 + q_t^2}$. Substituting $\vartheta = \theta_1 - \phi$, one gets

$$\int \frac{e}{2\pi} \begin{pmatrix} \sin \theta_1 \cos \vartheta - \cos \theta_1 \sin \vartheta \\ -\sin \theta_1 \sin \vartheta - \cos \theta_1 \cos \vartheta \end{pmatrix} e^{iq_1 R \cos \vartheta} d\vartheta = -ieJ_1(q_1 R) \begin{pmatrix} \sin \theta_1 \\ -\cos \theta_1 \end{pmatrix}, \quad (23)$$

Therefore, the momentum representation of the current is as follows:

$$j^\mu(q_1) = -\frac{ie}{q_1} J_1(q_1 R) \epsilon^{\mu\nu} q_1^\nu, \quad (24)$$

In the same way,

$$\int \frac{e}{2\pi R} \begin{pmatrix} -x \\ t \end{pmatrix} \frac{\delta(\sqrt{x^2 + t^2} - R)}{\sqrt{x^2 + t^2}} e^{i(\vec{q}_1 + \vec{q}_2 + \vec{q}_3, \vec{x})} dx dt = -ie J_1(q_4 R) \begin{pmatrix} \cos \theta_4 \\ -\sin \theta_4 \end{pmatrix}, \quad (25)$$

Therefore in the case of two particles the polarization operator in the momentum representation is

$$\pi^{\mu\nu}(q) = \frac{\Gamma}{2V} (e J_1(q R))^2 \left(\delta^{\mu\nu} - \frac{q^\mu q^\nu}{q^2} \right), \quad (26)$$

Here $\mu, \nu = 1, 2$. That coincides with the result [19]. In the four-particle case,

$$\pi^{\mu\nu\kappa\rho}(q_1, q_2, q_3, q_4) = \frac{\Gamma}{2V} \left(\prod \frac{e J_1(q_i R)}{q_i} \right) (q_1^\mu q_2^\nu - \delta^{\mu\nu}(q_1 q_2)) (q_3^\kappa q_4^\rho - \delta^{\kappa\rho}(q_3 q_4)). \quad (27)$$

with the constraint $\vec{q}_1 + \vec{q}_2 + \vec{q}_3 + \vec{q}_4 = 0$. Like in the scalar case, the result (27) can be generalized to the arbitrary number of the interacting particles.

All the previous calculations were carried out under the assumption of small momenta $q \ll m$. This means that the interaction between a photon and the wall of the bounce doesn't affect the shape of the latter. Generally speaking, the interaction with the photons considerably changes the shape of the bounce, and therefore changes the action. The modification of the action leads to an exponential correction to the rates of the processes. However, such modification of the action leads to significant complication of the setting, and this problem lies beyond the scope of the current paper.

In the next subsection the corrections because of the self-interaction of the bounce are considered. The model with the small momenta is showed to be accurate enough to obtain these exponential corrections for the electrodynamics and the scalar theory.

2.3 Coulomb and Yukawa corrections to the action

In the paper [9] the Coulomb correction to the action for the bounce was found. The action has the form

$$S = \frac{\pi m^2}{eE} - \frac{e^2}{4}, \quad (28)$$

Note that the Coulomb interaction doesn't affect either the shape or the size of the bounce. The width of the process acquires exponential correction, although small for weak fields.

$$\Gamma = \frac{(eE)^2}{(2\pi)^3} \exp \left(-\frac{\pi m^2}{eE} + \frac{e^2}{4} \right), \quad (29)$$




Figure 2: To obtain the Coulomb correction for the action of the bounce, the interaction with arbitrarily many photons is to be studied.

The correction to the width (29) implies the existence of the upper critical field. That is, when the field is $E_c \gtrsim \frac{m^2}{e^3}$, the process ceases to be exponentially suppressed. Similar phenomena are discussed in QCD [21].

The result (26) allows to obtain this correction in quite a simple fashion. Furthermore, the result (18) can be used to calculate the Yukawa correction for the scalar case, where the following on the [9] paper is troublesome.

To calculate the Coulomb correction, the interaction with a large number of photons is to be taken into account (see fig. 2). Let's calculate first a one-photon contribution. To do so, we have to integrate the operator (26) with the propagator of the photon.

$$\Gamma_1 = \frac{1}{2} \frac{\Gamma}{2V} \int_0^\infty (eJ_1(qR))^2 \left(\delta^{\mu\nu} - \frac{q^\mu q^\nu}{q^2} \right) \frac{\delta^{\mu\nu}}{q^2} q dq = \frac{\Gamma}{2V} \frac{e^2}{4}, \quad (30)$$

Note that this result doesn't depend on the radius of the bounce too. To compute the Coulomb correction, we have to sum up the contributions of any number of photons, taking into account the combinatorial factors.

$$\Gamma_C = \sum \frac{1}{i!} \Gamma_i = \frac{\Gamma}{2V} \exp \left(\frac{e^2}{4} \right), \quad (31)$$

So the correction to the action coincides with the result of [9]. As we know that in the momentum representation the polarization operator is a tensor product of some vectors, we can see that the Coulomb correction to the polarization operator is just the same.

$$\pi_C^{\mu\nu} = \pi^{\mu\nu} \exp \left(\frac{e^2}{4} \right). \quad (32)$$

The calculation (30) contains a certain subtlety. All the previous calculations are valid in the case of soft photons, $q \ll m$. In the area where the momentum of the photon is




Figure 3: The bounce in the case of hard photons. The shape significantly changes.

significant, the shape of the bounce ceases to be circular. The change of the form of the bounce leads to a correction of the action, and therefore to an exponential correction of the width (see fig. 3).

In the equation (30) we have integrated over all possible values of momentum; however, the integrand significantly differs from zero only in the area where $qR \sim 1$. Thus the correction of the shape is to be neglected.

Yukawa correction can be found in the same way. The width with one-particle correction is as following

$$\Gamma_1 = \frac{1}{2L} \frac{\Gamma}{2L} \int_0^\infty g^2 J_0(qR)^2 \frac{q dq}{q^2 + m^2} = \frac{\Gamma}{2L} \frac{g^2}{2} I_0(mR) K_0(mR), \quad (33)$$

The corrected width then is

$$\Gamma_Y = \frac{\varepsilon}{4\pi} \exp \left(-\frac{\pi\mu^2}{\varepsilon} + \frac{g^2}{2} I_0 \left(\frac{m\mu}{\varepsilon} \right) K_0 \left(\frac{m\mu}{\varepsilon} \right) \right). \quad (34)$$

This exponential correction $I_0 \left(\frac{m\mu}{\varepsilon} \right) K_0 \left(\frac{m\mu}{\varepsilon} \right)$ behaves like a logarithm for small argument. However, this is not an unusual property for a two-dimensional theory to have infrared logarithmic divergences. Bessel functions often appear in answers for two dimensions (see e.g. [16]).

This means that there exists some *lower* critical mass of the χ particle under which the process is no longer exponential suppressed.

In the scalar case the exponential correction due to the distortion of the shape of the bounce is also possible, therefore the area $m \ll \mu$ is considered. The lower critical mass lies in this area. Furthermore, the correction obtained grows small in the case $m > \mu$, that is, the main source of the correction in the case of massive χ particles is the distortion of the shape of the bounce.

Of course, the same correction factor arises in the cross-section and the n -fold correlator as well as in the width.

3 Conclusions

In this paper the two-photon-stimulated Schwinger process was considered by purely semiclassical means. We calculated the cross-section for two-photon interaction as leading term in $\frac{eE}{m^2}$ expansion which is applicable for the strength of the field much smaller than E_{crit} . This cross-section was studied in case of soft photons ($\omega \ll m$). In this case the shape of the bounce remains circular, and this fact allows to apply the procedure of calculating the imaginary part of a correlator to get the answer.

The consideration of the correlator appears to be simple and useful. The correlators take especially transparent form in the momentum representation, where they are just a product of the similar multiples. This makes possible the computation of the imaginary part of the correlator for the arbitrarily many interacting particles. The correction for the self-interaction of the bounce is also made, and not only in the Coulomb case, where the classical calculation (see [9]) is quite straightforward, but also in the Yukawa case, where the classical calculation involves a solution to a transcendent equation.

The authors are especially grateful to A. Gorsky for initiating the work and for useful and productive discussions. The work was supported by grants RFBR 09-02-00308 and FASI RF 14.740.11.0347.

References

- [1] J. Schwinger, Phys. Rev. **82**, 664 (1951).
- [2] W. Heisenberg, H. Euler, Z. Phys. **98** (1936) 714.
- [3] R. Schutzhold, H. Gies, G. Dunne, Phys. Rev. Lett. **101**, 130404 (2008), arXiv:0807.0754 [hep-th].
- [4] G. Dunne, H. Gies, R. Schutzhold, Phys. Rev. D **80**, 11301 (2009), arXiv:0908.0948 [hep-ph].
- [5] M. B. Voloshin, I. Yu. Kobzarev, L. B. Okun, Yad. Fiz. **20**, 1229 (1974).
- [6] M. B. Voloshin, Yad. Fiz. **42**, 1017 (1985).
- [7] C. G. Callan, S. Coleman, Phys. Rev. D **16**, 1762, (1977).
- [8] S. Coleman, Phys. Rev. D **15**, 2929 (1977).
- [9] I. K. Affleck, O. Alvarez and N. S. Manton, Nucl. Phys. B **197**, 509 (1982).
- [10] K. B. Selivanov and M. B. Voloshin, JETP Lett. **42**, 422 (1985).
- [11] V. G. Kiselev, Phys. Rev. D **45**, 2929 (1992).
- [12] A. S. Gorsky, K. A. Saraikin, K. G. Selivanov, Nucl. Phys., B **628**, (2002), 270-294, hep-th/0507174.

- [13] A. S. Gorsky, K. G. Selivanov, Nucl. Phys., B **571**, (2000), 120-134, hep-th/9904041.
- [14] A. S. Gorsky, K. G. Selivanov, Phys.Lett. B **485** (2000) 271-277, arXiv:hep-th/0005066v2
- [15] A. K. Monin, JHEP **109** 0510 (2005), hep-th/0509047.
- [16] A. Gorsky, M. B. Voloshin, Phys. Rev. D **73** (2006), arXiv:hep-th/0511095v1
- [17] A. K. Monin, A. V. Zayakin, JETP Lett. **84** (2006) 5-10, arXiv:hep-ph/0605079v2.
- [18] A. K. Monin, A. V. Zayakin, Phys.Rev.D **75**:065029,2007, arXiv:hep-ph/0611038v2.
- [19] A. Monin and M. B. Voloshin, Phys.Rev.D **81**:085014,2010, arXiv:1001.3354 [hep-th].
- [20] A. Monin and M. B. Voloshin, Phys.Rev.D **81**:025001,2010, arXiv:1004.2015 [hep-th].
- [21] G. Semenoff, K. Zarembo, arXiv:1109.2920v1 [hep-th].